

ON THE NATURALITY OF THE MATHAI-QUILLEN FORMULA

GYULA LAKOS

ABSTRACT. We give an alternative proof for the Mathai-Quillen formula for a Thom form using its natural behaviour with respect to fiberwise integration. We also study this phenomenon in general context.

1. THE MATHAI-QUILLEN FORMULA

Let $\pi : V \rightarrow M$ be an oriented vector bundle. A smooth, closed differential form Φ on V is called a Thom-form for π , if it is of vertically compact support, and its vertical integral $\pi_*\Phi = 1$. (See Bott and Tu, [2] for more details.) Nevertheless, it is enough to assume that Φ is only vertically rapidly decreasing, because by a simple push-forward operation one can bring it to vertically compact form. In what follows we mean rapidly decreasing Thom forms.

If $\pi : V \rightarrow M$ is an oriented bundle of rank n endowed with a metric then local trivializations can be chosen so that $x = [x_i]$ are the coordinates in the trivialized bundle, $dx_1 \wedge \dots \wedge dx_n$ is the orientation form of the bundle, and $x^2 = x^\top x = \sum x_i^2$ is the metric. Such trivializations are called compatible. If ∇ is a connection on the bundle then $\theta = [\theta_{ij}]$ are the connection 1-forms, assuming that ∇ is of local form $d + \theta \cdot$ as a covariant differential with respect to the coordinates x ; and $\Omega = d\theta + \theta\theta = [\Omega_{ij}]$ are the corresponding local curvature 2-forms. (If u is a column vector then we write $u_i = \mathbf{e}_i^\top u$ for its i -th element, and if M is a matrix, then we write $M_{ij} = \mathbf{e}_i^\top M \mathbf{e}_j$ for its (i, j) -th entry.)

The famous result of Mathai and Quillen in its most naive form says the following:

Theorem 1 (Mathai, Quillen, [3]). *If $\pi : V \rightarrow M$ is an oriented bundle of rank n endowed with a metric and a compatible connection ∇ then there is a well-defined Thom form $\text{TMQ}(V, \nabla)$ on V such that*

$$\text{TMQ}(V, \nabla) = \pi^{-n/2} e^{-x^2} \sum_{\substack{I \cup I' = \{1, \dots, n\}, \\ |I'| \text{ is even}}} (-1)^{(I, I')} (dx + \theta x)_{i_1} \dots (dx + \theta x)_{i_{|I|}} \\ \frac{1}{2^{|I'|} (|I'|/2)!} \sum_{\sigma \in \Sigma_{|I'|}} (-1)^\sigma \Omega_{i'_{\sigma(1)}, i'_{\sigma(2)}} \dots \Omega_{i'_{\sigma(|I'|-1)}, i'_{\sigma(|I'|)}}$$

in any compatible trivialization. (Here the convention is that $I = \{i_1, \dots, i_{|I|}\}$ with elements listed in increasing order; similarly for I' ; and (I, I') is the permutation which puts the elements of I in front of the elements of I' .)

The form $\text{TMQ}(V, \nabla)$ is rather explicit; the main difficulty lies in proving that it is well-defined and closed, even if it is a local computation. One quite simple

2000 *Mathematics Subject Classification.* Primary 58A10, Secondary 53C05, 57R20.

Key words and phrases. Mathai-Quillen formula, Thom form, Gaussian integrals, Wick formula.

proof can be found in the book [1] of Berline, Getzler, and Vergne using elementary supercalculus.

We give an alternative proof for Theorem 1 above, realizing a different strategy: One classical existence proof for the Thom form goes as follows: We embed V into a trivial bundle like

$$\begin{array}{ccc} V & \xleftarrow{\pi_1} & V \oplus W \simeq M \times \mathbb{R}^{n+m} \\ & \searrow \pi & \downarrow \\ & & M \end{array}$$

we take a Thom-form Φ_0 on the trivial bundle, and then we can take the fiberwise integral $\Phi = (\pi_1)_* \Phi_0$ as a Thom-form on V .

We will see that the Mathai-Quillen form occurs as the result of such a process. More specifically: A splitting of the trivial bundle $M \times \mathbb{R}^{n+m}$ can be realized by an involution valued $(n+m) \times (n+m)$ matrix function Q on M , where V is the (-1) -eigenspace of Q and W is the $(+1)$ -eigenspace. Let π_1 be the projection to the first factor. Let $y = [y_i]$ denote the column vector of the coordinates of the trivial bundle $M \times \mathbb{R}^{n+m}$. Now, there is a natural connection ∇_0 on the trivial bundle, such that in the coordinates y its covariant derivative is given as $d + \theta^{(y)} \cdot = d + \frac{1}{2} Q dQ \cdot$. This connection leaves the bundles V, W invariant. Furthermore, if we assume that Q is orthogonal then it is compatible to the metric which is the restriction of the metric y^2 from the trivial bundle. We denote the restriction of the connection above to V by $\nabla_0|_V$. Let us take the Thom form $\Phi_0 = \pi^{-(n+m)/2} e^{-y^2} dy_1 \wedge \dots \wedge dy_{n+m}$. Then the key statement is:

Theorem 2. *If V is oriented then*

$$\text{TMQ}(V, \nabla_0|_V) = (\pi_1)_* \Phi_0,$$

where $(\pi_1)_*$ is according to the complementary orientation on W .

(The statement includes the well-definedness of the left side.)

One can ask that how general the pairs $(V, \nabla_0|_V)$ are among arbitrary metric bundle and connection pairs. The answer is that they are completely general: According to a theorem of Narasimhan and Ramanan [4], every metric pair can be realized like that. (Their formalism is a bit different.) As a corollary we have Theorem 1. In fact, we are interested in a local computation, so we do not even have to use [4] in its full power.

Proof of Theorem 2. Locally, we can always take an $\text{SO}(n+m)$ -valued function A such that $Q = A Q_0 A^{-1}$, where $Q_0 = \begin{bmatrix} -1_{n \times n} & \\ & 1_{m \times m} \end{bmatrix}$ is a fixed matrix. Let us then define $x = A^{-1}y$ as the new set of coordinates. We see that y is in the ± 1 -eigenspace of Q if and only if x is in the ± 1 -eigenspace of Q_0 . From this it follows that the coordinates $x_1, \dots, x_n$ trivialize V . Actually more is true: $x_{n+1}, \dots, x_{n+m}$ trivialize W . This is more than we expect (we want to trivialize V only), nevertheless we can note that these trivializations account for all possible compatible trivializations of V with respect to the metric, because such a local trivialization of V can always be extended to a well-oriented orthogonal trivialization of $V \oplus W$. It is practical to incorporate the variables $x_1, \dots, x_n$ into a column vector x_o , and the variables $x_{n+1}, \dots, x_{n+m}$ into the column vector x_h . With some abuse of notation $x = x_o + x_h$.

Now, we have a connection on $M \times \mathbb{R}^{n+m}$, whose covariant derivative can be written as $d + \theta^{(y)} \cdot = d + \frac{1}{2} Q dQ \cdot$ with respect to the coordinate functions y . Then

the covariant derivative is given as

$$d + \theta^{(x)} \cdot = (A^{-1} \cdot) (d + \frac{1}{2} Q d Q \cdot) (A \cdot) = d + \frac{1}{2} (A^{-1} d A + Q_0 A^{-1} d A Q_0) \cdot$$

with respect to the coordinate functions x . The curvature form is $\Omega^{(y)} = \frac{1}{4} d Q d Q$ with respect to y . Similarly, with respect to x it is given as

$$\Omega^{(x)} = -\frac{1}{4} (A^{-1} d A - Q_0 A^{-1} d A Q_0) (A^{-1} d A - Q_0 A^{-1} d A Q_0).$$

Using the block decomposition $A^{-1} d A = \begin{bmatrix} (A^{-1} d A)_{oo} & (A^{-1} d A)_{oh} \\ (A^{-1} d A)_{ho} & (A^{-1} d A)_{hh} \end{bmatrix}$ we find

$$\theta^{(x)} = \begin{bmatrix} (A^{-1} d A)_{oo} & \\ & (A^{-1} d A)_{hh} \end{bmatrix},$$

$$\Omega^{(x)} = - \begin{bmatrix} (A^{-1} d A)_{oh} (A^{-1} d A)_{ho} & \\ & (A^{-1} d A)_{ho} (A^{-1} d A)_{oh} \end{bmatrix}.$$

In particular, restricted to V , with respect to the coordinate functions x_o , it yields

$$(\dagger) \quad \theta = (A^{-1} d A)_{oo}, \quad \text{and} \quad \Omega = -(A^{-1} d A)_{oh} (A^{-1} d A)_{ho}.$$

Let us note that Φ_0 can be written as $\pi^{-(n+m)/2} e^{-y^2} (A^{-1} dy)_1 \wedge \dots \wedge (A^{-1} dy)_{n+m}$, because A is $\text{SO}(n+m)$ -valued. Applying $y = Ax$ we find that

$$\Phi_0 = \pi^{-(n+m)/2} e^{-x^2} (dx + A^{-1} d A x)_1 \wedge \dots \wedge (dx + A^{-1} d A x)_{n+m}.$$

In this local coordinate system the fiberwise integration $(\pi_1)_*$ is just integration in the variables $x_{n+1}, \dots, x_{n+m}$ whose infinitesimal forms appear only in the last m terms in the integrand. So, it yields

$$(\pi_1)_* \Phi = \pi^{-(n+m)/2} \int e^{-x^2} (dx + A^{-1} d A x)_1 \wedge \dots \wedge (dx + A^{-1} d A x)_n dx_{n+1} \dots dx_{n+m}.$$

Let us decompose the column vector x as $x_o + x_h$. Then x_o contains the non-integration variables $x_1, \dots, x_n$, while x_h contains the integration variables $x_{n+1}, \dots, x_{n+m}$. The term $(dx + A^{-1} d A x)_i$ decomposes as $dx_i + ((A^{-1} d A)_{oo} x_o)_i + ((A^{-1} d A)_{oh} x_h)_i$. After reordering we find that $(\pi_1)_* \Phi$

$$= \pi^{-(n+m)/2} e^{-x_o^2} \sum_{I \cup I' = \{1, \dots, n\}} (dx_o + (A^{-1} d A)_{oo} x_o)_{i_1} \dots (dx_o + (A^{-1} d A)_{oo} x_o)_{i_{|I|}}$$

$$(-1)^{(I, I')} \int e^{-x_h^2} ((A^{-1} d A)_{oh} x_h)_{i'_1} \dots ((A^{-1} d A)_{oh} x_h)_{i'_{|I'|}} dx_{n+1} \dots dx_{n+m}$$

(the sign change is due to the fact that 1-forms anticommute).

Let us remind ourselves to the Wick formula of Gaussian integrals. It says that if we are in $\mathbb{R}^l$, $z = [z_k]$ is the column vector of the variables, $b_1, \dots, b_s$ are scalar-valued fixed column vectors then

$$\int e^{-z^2} (b_1^\top z) \dots (b_s^\top z) = \frac{\pi^{l/2}}{2^s (s/2)!} \sum_{\sigma \in \Sigma_s} (b_{\sigma(1)}^\top b_{\sigma(2)}) \dots (b_{\sigma(l-1)}^\top b_{\sigma(l)}),$$

if s is even, and $= 0$ if s is odd. If the column vectors b_i are 1-form valued then the expression remains the same except that $(-1)^\sigma$ must be inserted after $\sum$, because interchanging the order of the coefficients yields some sign changes.

Applying the formula with respect to $m, x_h, \mathbf{e}_{i_j}^\top (A^{-1}dA)_{oh}$ in the place of $l, z, b_j^\top$, respectively, we find

$$\begin{aligned} (\pi_1)_* \Phi_0 &= \pi^{-n/2} e^{-x_o^2} \sum_{\substack{I \cup I' = \{1, \dots, n\}, \\ |I'| \text{ is even}}} (-1)^{(I, I')} (dx_o + (A^{-1}dA)_{oo} x_o)_{i_1} \dots \\ &\quad \dots (dx_o + (A^{-1}dA)_{oo} x_o)_{i_{|I|}} \frac{1}{2^{|I'|} (|I'|/2)!} \sum_{\sigma \in \Sigma_{|I'|}} (-1)^\sigma \\ &\quad ((A^{-1}dA)_{oh} (A^{-1}dA)_{oh}^\top)_{i'_{\sigma(1)}, i'_{\sigma(2)}} \dots ((A^{-1}dA)_{oh} (A^{-1}dA)_{oh}^\top)_{i'_{\sigma(|I'|-1)}, i'_{\sigma(|I'|)}}. \end{aligned}$$

From orthogonality $(A^{-1}dA)_{oh}^\top = -(A^{-1}dA)_{ho}$, and applying $(\ddagger)$ we obtain

$$\begin{aligned} (\pi_1)_* \Phi_0 &= \pi^{-n/2} e^{-x_o^2} \sum_{\substack{I \cup I' = \{1, \dots, n\}, \\ |I'| \text{ is even}}} (-1)^{(I, I')} (dx_o + \theta x_o)_{i_1} \dots (dx_o + \theta x_o)_{i_{|I|}} \\ &\quad \frac{1}{2^{|I'|} (|I'|/2)!} \sum_{\sigma \in \Sigma_{|I'|}} (-1)^\sigma \Omega_{i'_{\sigma(1)}, i'_{\sigma(2)}} \dots \Omega_{i'_{\sigma(|I'|-1)}, i'_{\sigma(|I'|)}}. \end{aligned}$$

The formula applies to every compatible trivialization x_o of V locally. This implies that the Mathai-Quillen form is well-defined on V , and Theorem 2 holds. $\square$

2. CONSTRAINED CONNECTIONS AND FIBERWISE INTEGRALS

We can put Theorem 2 into more general context. In general, assume that ∇ is a connection on a bundle $\mathcal{V}$, and Q is an involution-valued function on the bundle, the subbundles V and W are the (-1) - and $(+1)$ -eigenspaces of Q . Then one can define the constrained connection

$$\nabla_Q = \frac{1}{2}(\nabla + (Q \cdot) \nabla(Q \cdot)) = \nabla + \frac{1}{2}Q(\nabla Q) \cdot \quad .$$

This connection leaves the bundles V and W invariant; in particular, $\nabla_Q|_V$ can be taken. Now, if $\mathcal{V}$ is a metric bundle, and V is a subbundle then it is reasonable to define the constrained connection $\nabla|_V$ as $\nabla_Q|_V$, where Q is that unique orthogonal involution-valued function whose (-1) -eigenspace is V . In this case one can simply describe the constrained connection $\nabla|_V$ locally as follows: Assume that the variables $x_1, \dots, x_n$ trivialize the bundle $\mathcal{V}$, and $x_1, \dots, x_k$ trivialize V , and the metric is given by x^2 . Assume that the connection ∇ as a covariant derivative in the coordinates x is given by the connection form θ . Then we can just take that covariant derivative on W whose connection form is just $\theta|_{k \times k}$, ie. θ restricted to the variables $x_1, \dots, x_k$. This constrain operation has many natural properties, which are immediate to see, for example:

Lemma 1. (a) If $V_2 \subset V_1 \subset \mathcal{V}$ then $(\nabla|_{V_1})|_{V_2} = \nabla|_{V_2}$.

(b) If $V_1 \subset \mathcal{V}_1$, $V_2 \subset \mathcal{V}_2$ then $(\nabla_1 \oplus \nabla_2)|_{V_1 \oplus V_2} = \nabla_1|_{V_1} \oplus \nabla_2|_{V_2}$. $\square$

Let us take the orthogonal decomposition $V \oplus V^\top = \mathcal{V}$, and the projection π_1 to the first factor.

Theorem 3. If $\mathcal{V}$ and V are oriented then

$$\text{TMQ}(V, \nabla|_V) = (\pi_1)_* \text{TMQ}(\mathcal{V}, \nabla)$$

where $(\pi_1)_*$ is according to the complementary orientation on $V^\top$.

Clearly, Theorem 2 is the special case when the pair $(\mathcal{V}, \nabla)$ is trivial.

Proof. According to [4] locally we can embed $(\mathcal{V}, \nabla)$ into the trivial pair $(M \times \mathbb{R}^{n+m}, \nabla_0)$ such that $\nabla = \nabla_0|_V$. Then Lemma 1.a, and the fact that fiberwise integration has the similar property implies our statement through Theorem 2. $\square$

We remark that Theorem 3 is not hard to prove with simple supercalculus directly; the calculations are similar to the ones in the proof of Theorem 2.

Using elementary properties of the constrained connections we can also recover the behaviour of the Mathai-Quillen form with respect to patchings.

Assume that $(\mathcal{V}_1, \nabla_1), \dots, (\mathcal{V}_s, \nabla_s)$ are metric bundle and connection pairs. Let $(\mathcal{V}, \nabla)$ be the direct sum of these pairs. Assume that V is a metric vector bundle and $\iota_s : V \rightarrow \mathcal{V}_s$ are metric inclusions of bundles. Assume that $(\xi_1, \dots, \xi_s)$ is a quadratic partition of unity, ie. $\sum \xi_s^2 = 1$. Let $\iota : V \rightarrow \mathcal{V}$ be the inclusion given by

$$\iota(x) = \xi_1 \iota_1(x) + \dots + \xi_s \iota_s(x).$$

Then we claim:

Lemma 2.

$$\iota^*(\nabla|_{\iota(V)}) = \xi_1^2 \iota_1^*(\nabla_1|_{\iota_1(V)}) + \dots + \xi_s^2 \iota_s^*(\nabla_s|_{\iota_s(V)}).$$

Proof. By [4], Lemma 1, and locality, it is enough to prove the statement when V is $M \times \mathbb{R}^n$, and $\mathcal{V}_i$ are copies of $M \times \mathbb{R}^{n+m}$ with trivial connections. Locally we can extend the inclusions $\iota_k : M \times \mathbb{R}^n \rightarrow M \times \mathbb{R}^{n+m}$ to metric isomorphisms $A_k : M \times \mathbb{R}^{n+m} \rightarrow M \times \mathbb{R}^{n+m}$. Let us consider the following block matrix, where every block is an $(n+m) \times (n+m)$ matrix:

$$A = \begin{bmatrix} 0 & -\xi_1 \text{Id} & -\xi_2 \text{Id} & \dots & -\xi_s \text{Id} \\ \xi_1 A_1 & (1 - \xi_1^2) A_1 & -\xi_1 \xi_2 A_1 & \dots & -\xi_1 \xi_s A_1 \\ \xi_2 A_2 & -\xi_2 \xi_1 A_2 & (1 - \xi_2^2) A_2 & & -\xi_2 \xi_s A_2 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ \xi_s A_s & -\xi_s \xi_1 A_s & -\xi_s \xi_2 A_s & \dots & (1 - \xi_s^2) A_s \end{bmatrix}.$$

One can see that A is an orthogonal matrix. Let Q_0 be that matrix which is the $(n+m)(s+1) \times (n+m)(s+1)$ identity matrix, except that its first n diagonal entries are -1 . Let X_{oo} denote the restriction of a matrix X to its top left $n \times n$ submatrix (for block matrices they fall into the first block). Then a straightforward matrix computation yields:

$$(A^{-1} dA)_{oo} = \sum_{i=1}^s \xi_i^2 (A_i^{-1} dA_i)_{oo}.$$

According to the argument in the proof of Theorem 2, which culminated in $(\ddagger)$, we see that the meaning of this equation is that the (-1) -eigenspace bundle of Q_0 pushed forward by A obtains a connection through the constrain operation such that its pull-back through A is the linear combination of the similar connections obtained through A_i , which fact in the present situation is equivalent to the formula of the statement. $\square$

In fact, we may notice that the explicit part of the calculation above is essentially the same one which makes possible the result of [4]. As a corollary we obtain:

Theorem 4. *Let V be a metric bundle and $\nabla_1, \dots, \nabla_s$ be various compatible connections. Let $\xi_1, \dots, \xi_s$ be a quadratic partition of unity. Let the inclusion $\iota : V \rightarrow V \oplus \dots \oplus V$ be defined as before. Then*

$$\text{TMQ}(V, \xi_1^2 \nabla_1 + \dots + \xi_s^2 \nabla_s) = \iota^*(\pi_\xi)_* \text{TMQ}(V \oplus \dots \oplus V, \nabla_1 \oplus \dots \oplus \nabla_s),$$

where π_ξ is the orthogonal projection to $\iota(V)$ given by the matrix $[\xi_i \xi_j \text{Id}_V]_{ij}$.

Proof. This follows from Lemma 2 and Theorem 3, when the inclusions ι_k are isomorphisms. $\square$

REFERENCES

- [1] N. Berline, E. Getzler, M. Vergne: *Heat kernels and Dirac operators*. 2nd ed. Grundlehren der mathematischen Wissenschaften 298, Springer-Verlag, Berlin, 1998.
- [2] R. Bott, L. W. Tu: *Differential forms in algebraic topology*. Graduate Texts in Mathematics 82, Springer-Verlag, Berlin, 1982.
- [3] V. Mathai, D. Quillen: Superconnections, Thom classes and equivariant differential forms. *Topology* **25** (1986), 85–110.
- [4] M. S. Narasimhan, S. Ramanan: Existence of universal connections. *Amer. J. Math.* **83** (1961), 583–572.

DEPARTMENT OF GEOMETRY, EÖTVÖS UNIVERSITY, PÁZMÁNY PÉTER S. 1/C, BUDAPEST, H-1117, HUNGARY

E-mail address: lakos@cs.elte.hu